



LETTER TO EDITOR

Potential pathogenesis of ageusia and anosmia in COVID-19 patients.

What we know from the literature

Luigi Angelo Vaira^{1,*} MD; Giovanni Salzano² MD; Alessandro Giuseppe Fois³ MD;
Pasquale Piombino² MD, FEBOMFS; Giacomo De Riu¹MD, FEBOMFS.

¹ Maxillofacial Surgery Unit, University Hospital of Sassari, Viale San Pietro 43/B, 07100 Sassari, Italy.

² Maxillofacial Surgery Unit, University Hospital of Naples “Federico II”, Via Pansini 5, 80131 Naples, Italy.

³ Respiratory Diseases Operative Unit, University Hospital of Sassari, Viale San Pietro 43/B, 07100 Sassari, Italy.

*Corresponding Author:

Luigi Angelo Vaira

Viale San Pietro 43/B

e-mail: luigi.vaira@gmail.com

tel.: +393401846168

fax: +39079229002

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/alr.22593](https://doi.org/10.1111/alr.22593).

This article is protected by copyright. All rights reserved.

Keywords: Covid-19; ageusia; anosmia; smell disorders; taste disorders; coronavirus; SARS-CoV-2

Abstract

From the first reports, ageusia and anosmia appear to be frequent clinical features in coronavirus disease 19 (COVID-19) patients. We have performed a survey of the literature, analyzing the possible causes of these chemosensory alterations, which may be useful as a starting point for specific further studies.

Dear Editor,

The rapid spread of the Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) in Europe and America is enabling researchers to acquire a large amount of clinical data. Interestingly, some of these findings differ significantly from those reported in Chinese case studies. In particular, the presence of taste and smell alterations seems to be a frequent clinical feature of the coronavirus disease 19 (COVID-19), with a frequency ranging from 19.4% to 88% of patients¹⁻⁴.

The exact pathogenesis of these chemosensitive disorders has not yet been clarified. At the same time, the comprehension of these mechanisms could be fruitful to understand the way in which the virus spreads through the organism.

We report our survey of the literature up to April 14, 2020, analyzing the possible causes of these chemosensory alterations, which may be useful as a starting point for specific future studies.

Ageusia

There are no studies in the literature regarding the possible relationship between coronavirus and the development of taste disorders. Angiotensin-converting enzyme 2 (ACE2) has been identified as the cellular receptor for SARS-CoV-2⁵. ACE2 receptors are expressed diffusely on the mucous membrane of the whole oral cavity, particularly on the tongue⁶. The role of ACE2 in modulating taste perception has been highlighted in many studies analyzing the chemosensitive side effects of ACE2 inhibitors and angiotensin II blockers^{7,8}. The mechanism by which ACE2 inhibitors cause taste disturbance is unclear but does not appear to be related to any alteration in serum^{7,8} and saliva⁷ zinc levels. Presumably, these drugs inactivate the G-protein-coupled proteins and sodium-ion channels present in the taste receptors⁵. The taste disturbance generally regresses after the suspension of treatment⁸.

Moreover, the Middle East Respiratory Syndrome (MERS) coronavirus may bind to the sialic acid receptors⁹, an ability which has also recently been described for SARS-CoV-2¹⁰. Sialic acid is a fundamental component of the salivary mucin, and it protects the glycoproteins that convey gustatory molecules inside the taste pores from premature enzymatic degradation¹¹. A reduction of sialic acid in the saliva is associated with an increase in the gustatory threshold¹². In such a way, SARS-CoV-2 could therefore occupy the binding sites of sialic acid on the taste buds, accelerating the degradation of the gustatory particles.

Another possibility is that the ability to perceive flavors in these patients is adversely affected by the concomitant presence of olfactory disturbances, due to the intimate functional correlation between these two chemosensory systems¹³. However, in the first published case series, taste dysfunctions are always more frequent than olfactory disturbances, presenting alone in 10.2-22.5% of patients²⁻⁴. This finding could lead to the hypothesis that the correlation between the two dysfunctions is not so close and that there are other factors behind the gustatory disorders in these patients.

Anosmia

ACE2 is expressed on the nasal mucosa, where it participates in respiratory inflammatory diseases by regulating the levels of inflammatory peptides, such as bradykinin¹⁴. However, in COVID-19 patients there does not seem to be present such an important inflammatory component and the alteration of the sense of smell is generally not accompanied by rhinitis symptoms^{1,3}.

Therefore, one hypothesis could be that the alterations are due to damage caused by the virus to the olfactory pathways. Anosmia has already been described in common coronavirus infections^{15,16}. In 2006, Hwang¹⁷ described a case of anosmia that persisted for two years after SARS had been contracted. In 2001, Schwob et al.¹⁸ and Youngentob et al.¹⁹ evaluated the anatomopathological and functional consequences of olfactory bulb line variant mouse hepatitis virus intranasal inoculation in murine models. The histopathological analysis of the anosmic mice revealed a minimal destruction of the olfactory epithelium which was nonetheless abnormal due to the predominance of immature neurons, an indication of an accelerated cell turnover^{18,19}. The reduction in the neuronal lifespan in the epithelium is most likely to be the consequence of a decrease in the trophic support supplied by the bulb to the

sensory neurons, caused by the loss of mitral cells and the absence of dendrites of the surviving mitral and tufted cells from the glomerular layer^{18,19}. Accordingly, the virus may carry out its damage at the level of the olfactory bulb rather than the epithelium. However, the coronavirus used in these studies has been laboratory modified to target specifically the olfactory bulb. This is not necessarily the same for SARS-CoV-2, which we know has different organ targets.

Nevertheless, these authors, while clarifying the target of the virus, did not provide clear indications on the exact pathogenesis of neuronal death. In 2008, Netland et al.²⁰ reported the effects of the SARS-CoV infection on the central central nervous system in mice transgenic for the human ACE2 receptor. The authors, not detecting any signs of inflammatory infiltration, hypothesized that neuronal death occurs due to a storm of cytokines, in particular IL-6, produced by neurons under the stimulation of the viral N-spike²⁰.

However, recent COVID-19 case series reports are showing a high rate of recovery of olfactory function within 1-2 weeks after the onset of the dysfunction^{3,4,21}. Furthermore, it would seem that the frequency (around 25%)²² of central nervous system symptoms is much lower than that of olfactory disorders. For these reasons, it is reasonable to hypothesize that the olfactory disorders are not related to, directly or indirectly, definitive viral damage to the neuronal cells. Conversely, the target of the virus may not be the neurons but other non-neuronal cells that express ACE2 receptors such as the olfactory epithelium sustentacular cells, microvillar cells, Bowman's gland cells, horizontal basal cells and olfactory bulb pericytes²³. Brann et al.²³, speculated that the loss of smell reported by COVID-19 patients is due to the infection of the supporting cells and vascular pericytes of the olfactory epithelium and bulb which, consequently, alters the function of the olfactory neurons. The further

involvement of stem cells (which express lower levels of ACE2 receptors) could be the basis of the long-lasting olfactory dysfunctions²³.

Future research lines

In conclusion, from the first reports¹⁻⁴, ageusia and anosmia appear to be a frequent clinical feature during COVID-19. The pathogenic mechanism underlying the chemosensitive disorders in these patients has not yet been clarified. In our opinion, three lines of research could be useful to provide important indications regarding some still unclear pathogenic aspects. First, it will be necessary to obtain clinical data on large patient cohorts to determine the exact frequency of these symptoms and monitor their recovery over time. Secondly, it may be useful to evaluate the correlation between the viral load in the nasopharyngeal swab and the extent of these chemosensitive disorders. Concluding, the histopathological analysis on samples obtained from patients who have died due to COVID-19 will be crucial to understand the nature and location of the olfactory dysfunction.

DISCLOSURES

Competing interests: none

Sponsorships: none

Funding sources: none

ACNOWLEDGMENTS

None declared.

References

- [1] Vaira LA, Salzano G, Deiana G, De Riu G. Ageusia and anosmia: common findings in COVID-19 patients. *Laryngoscope*. 2020; Doi:10.1002/lary.28692.
- [2] Giacomelli A, Pezzati L, Conti F et al. Self-reported olfactory and taste disorders in SARS-CoV-2 patients: a cross-sectional study. *Clin Infect Dis*. 2020; Doi: 10.1002/jmv.25794.
- [3] Lechien JR, Chiesta-Estomba CM, De Siati DR, et al. Olfactory and gustatory dysfunctions as a clinical presentation of mild-to-moderate forms of the coronavirus disease (COVID-19): a multicenter European study. *Eur Arch Otorhinolaryngol*. 2020. Doi: 10.1007/s00405-020-05965-1.
- [4] Yan CH, Faraji F, Prajapati DP, Boone CE, DeConde AS. Association of chemosensory dysfunction and Covid-19 in patients presenting with influenza-like symptoms. *Int Forum Allergy Rhinol*. 2020. Doi: 10.1002/air.22579.
- [5] Zhou P, Yang X-L, Wang X-G et al. A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature*. 2020;579:270-73.
- [6] Xu H, Zhong L, Deng J et al. High expression of ACE2 receptor of 2019-nCoV on the epithelial cells of oral mucosa. *Int J Oral Sci*. 2020;12:8
- [7] Tsuruoka S, Wakaumi M, Araki N, Ioka T, Sugimoto K, Fujimura A. Comparative study of taste disturbance by losartan and perindopril in healthy volunteers. *J Clin Pharmacol*. 2005;45-1319-23.

- [8] Suliburska J, Duda G, Pupek-Musialik D. The influence of hypotensive drugs on the taste sensitivity in patients with primary hypertension. *Acta Pol Pharm.* 2012;60:121-7
- [9] Park YJ, Walls AC, Wang Z et al. Structures of MERS-CoV spike glycoprotein in complex with sialoside attachment receptors. *Nat Struct Mol Biol.* 2019;26:1151-7.
- [10] Milanetti M, Miotto M, Di Rienzo L, Monti M, Gosti G, Ruocco G. In-Silico evidence for two receptors based strategy of SARS-CoV-2. *ArXiv:* 2003.11107.
- [11] Witt M, Miller Jr IJ. Comparative lectin histochemistry on taste bud in foliate, circumvallate and fungiform papillae of the rabbit tongue. *Histocemistry.* 119;98:173-82.
- [12] Pushpass RG, Pellicciotta N, Kelly C, Proctor G, Carpenter GH. Reduced salivary mucin binding and glycosylation in older adults influences taste in an in vitro cell model. *Nutrients.* 2019;11:228
- [13] Small DM, Prescott J. Odor/taste integration and the perception of flavor. *Exp Brain Res.* 2005;166:345-57.
- [14] Ohkubo K, Lee CH, Baraniuk JN, Merida M, Hausfeld JN, Kaliner MA. Angiotensin-converting enzyme in the human nasal mucosa. *Am J Respir Cell Mol Biol.* 1994;11:173-80.
- [15] De Haro-Licer J, Roura-Moreno J, Vizitiu A, González-Fernández A, González-Ares JA. Long term serious olfactory loss in cold and/or flu. *Acta Otorinolaringol Esp.* 2013;64:331-8.
- [16] Suzuki M, Saito K, Min WP et al. Identification of viruses in patients with postviral olfactory dysfunction. *Laryngoscope.* 2007;117:272-7.
- [17] Hwang CS. Olfactory neuropathy in severe acute respiratory syndrome: report of a case. *Acta Neurol Taiwan.* 2006;15:26-8.

- [18] Schwob JE, Saha S, Youngentob SL, Burk J. Intranasal inoculation with olfactory bulb line variant of mouse hepatitis virus causes extensive destruction of the olfactory bulb and accelerated turnover of neurons in the olfactory epithelium of mice. *Chem Senses*. 2001;26:937-52.
- [19] Youngentob SL, Schwob JE, Saha S, Manglapus G, Jubelt B. Functional consequences following infection of the olfactory system by intranasal infusion of the olfactory bulb line variant (OBLV) of mouse hepatitis strain JHM. *Chem Senses*. 2001;26:953-63.
- [20] Netland J, Meyerholz DK, Moore S, Cassel M, Perlman S. Severe acute respiratory syndrome coronavirus infection causes neuronal death in the absence of encephalitis in mice transgenic for human ACE2. *J Virol*. 2008;82:7264:75.
- [21] Vaira LA, Deiana G, Fois AG et al. Objective evaluation of anosmia and ageusia in COVID-19 patients: a single-center experience on 72 cases. *Head Neck*. 2020. Doi: 10.1002/HED.26204
- [22] Mao L, Wang M, Chen S et al. Neurological manifestations of hospitalized patients with COVID-19 in Wuhan, China: a retrospective case series study. *MedRxiv*. Doi: <https://doi.org/10.1101/2020.02.22.20026500>.
- [23] Brann DH, Tsukahara T, Weinreb C et al. Non-neuronal expression of SARS-CoV-2 entry genes in the olfactory system suggests mechanisms underlying COVID-19-associated anosmia. *BioRxiv*. 2020. Doi: <https://doi.org/10.1101/2020.03.25.009084>